

Can agriculture reach net zero emissions by 2050?"

Louis Verchot
Alliance Bioversity-CIAT
CGIAR
l.verchot@cgiar.org

Conclusions

Significant GHG emissions reductions in agriculture are achievable through reducing emission intensity of production, but population growth and dietary changes may offset absolute emissions reductions.

Emission reductions consistent with 2030 targets are achievable in many countries; attaining the 2050 targets will require innovation and systems transformation.

Net zero agriculture cannot be achieved without sinks!

Governance, economics, and sociocultural factors are the keys to food system transformations.

Production is the major source of emissions in food systems



Mitigate+: Research
for Low-Emission
Food Systems




Table SPM1. Net anthropogenic emissions due to Agriculture, Forestry, and other Land Use (AFOLU) and non-AFOLU (Panel 1)																				
		Direct Anthropogenic																		
Gas	Units	Net anthropogenic emissions due to Agriculture, Forestry, and Other Land Use (AFOLU)				Non-AFOLU anthropogenic GHG emissions ⁶		Total net anthropogenic emissions (AFOLU + non-AFOLU) by gas			AFOLU as a % of total net anthropogenic emissions, by gas		Natural response of land to human-induced environmental change ⁷			Net land – atmosphere flux from all lands				
Panel 1: Contribution of AFOLU																				
		FOLU		Agriculture		Total														
		A		B		C = A + B		D		E = C + D		F = (C/E) *100			G			A + G		
CO ₂ ²																				
	Gt CO ₂ y ⁻¹	5.2 ± 2.6		No data ¹¹		5.2 ± 2.6		33.9 ± 1.8		39.1 ± 3.2		13%			-11.2	±	2.6	-6.0	±	3.7
CH ₄ ^{3,8}	Mt CH ₄ y ⁻¹	19.2 ± 5.8		141.6 ± 42.5		160.8 ± 43		201.3 ± 100.6		362 ± 109										
	Gt CO ₂ e y ⁻¹	0.5 ± 0.2		4.0 ± 1.2		4.5 ± 1.2		5.6 ± 2.8		10.1 ± 3.1		44%								
N ₂ O ^{3,8}	Mt N ₂ O y ⁻¹	0.3 ± 0.1		8.3 ± 2.5		8.7 ± 2.5		2.0 ± 1.0		10.6 ± 2.7										
	Gt CO ₂ e y ⁻¹	0.09 ± 0.03		2.2 ± 0.7		2.3 ± 0.7		0.5 ± 0.3		2.8 ± 0.7		81%								
Total (GHG)	Gt CO ₂ e y ⁻¹	5.8 ± 2.6		6.2 ± 1.4		12.0 ± 2.9		40.0 ± 3.4		52.0 ± 4.5		23%								

Table SPM1. Net anthropogenic emissions due to Agriculture, Forestry, and other Land Use (AFOLU) and non-AFOLU (Panel 1)

		Direct Anthropogenic																			
Gas	Units	Net anthropogenic emissions due to Agriculture, Forestry, and Other Land Use (AFOLU)					Non-AFOLU anthropogenic GHG emissions ⁶			Total net anthropogenic emissions (AFOLU + non-AFOLU) by gas			AFOLU as a % of total net anthropogenic emissions, by gas			Natural response of land to human-induced environmental change ⁷			Net land – atmosphere flux from all lands		
Panel 1: Contribution of AFOLU																					
		FOLU		Agriculture		Total															
		A		B		C = A + B		D		E = C + D			F = (C/E) *100			G			A + G		
CO ₂ ²																					
	Gt CO ₂ y ⁻¹	5.2	± 2.6	No data ¹¹		5.2	± 2.6	33.9	± 1.8	39.1	±	3.2	13%			-11.2	±	2.6	-6.0	±	3.7
CH ₄ ^{3,8}	Mt CH ₄ y ⁻¹	19.2	± 5.8	141.6	± 42.5	160.8	± 43	201.3	± 100.6	362	±	109									
	Gt CO ₂ e y ⁻¹	0.5	± 0.2	4.0	± 1.2	4.5	± 1.2	5.6	± 2.8	10.1	±	3.1	44%								
N ₂ O ^{3,8}	Mt N ₂ O y ⁻¹	0.3	± 0.1	8.3	± 2.5	8.7	± 2.5	2.0	± 1.0	10.6	±	2.7									
	Gt CO ₂ e y ⁻¹	0.09	± 0.03	2.2	± 0.7	2.3	± 0.7	0.5	± 0.3	2.8	±	0.7	81%								
Total (GHG)	Gt CO ₂ e y ⁻¹	5.8	± 2.6	6.2	± 1.4	12.0	± 2.9	40.0	± 3.4	52.0	±	4.5	23%								

Table SPM1. Net anthropogenic emissions due to Agriculture, Forestry, and other Land Use (AFOLU) and non-AFOLU (Panel 1)

		Direct Anthropogenic													
Gas	Units	Net anthropogenic emissions due to Agriculture, Forestry, and Other Land Use (AFOLU)			Non-AFOLU anthropogenic GHG emissions ⁶	Total net anthropogenic emissions (AFOLU + non-AFOLU) by gas		AFOLU as a % of total net anthropogenic emissions, by gas		Natural response of land to human-induced environmental change ⁷			Net land – atmosphere flux from all lands		
Panel 1: Contribution of AFOLU															
		FOLU		Agriculture	Total										
		A		B	C = A + B		D	E = C + D		F = (C/E) *100		G		A + G	
CO ₂ ²															
	Gt CO ₂ y ⁻¹	5.2 ± 2.6		No data ¹¹	5.2 ± 2.6		33.9 ± 1.8	39.1 ± 3.2		13%		-11.2	±	2.6	-6.0 ± 3.7
CH ₄ ^{3,8}	Mt CH ₄ y ⁻¹	19.2 ± 5.8		141.6 ± 42.5	160.8 ± 43		201.3 ± 100.6	362 ± 109							
	Gt CO ₂ e y ⁻¹	0.5 ± 0.2		4.0 ± 1.2	4.5 ± 1.2		5.6 ± 2.8	10.1 ± 3.1		44%					
N ₂ O ^{3,8}	Mt N ₂ O y ⁻¹	0.3 ± 0.1		8.3 ± 2.5	8.7 ± 2.5		2.0 ± 1.0	10.6 ± 2.7							
	Gt CO ₂ e y ⁻¹	0.09 ± 0.03		2.2 ± 0.7	2.3 ± 0.7		0.5 ± 0.3	2.8 ± 0.7		81%					
Total (GHG)	Gt CO ₂ e y ⁻¹	5.8 ± 2.6		6.2 ± 1.4	12.0 ± 2.9		40.0 ± 3.4	52.0 ± 4.5		23%					

Table SPM1. Net anthropogenic emissions due to Agriculture, Forestry, and other Land Use (AFOLU) and non-AFOLU (Panel 1)

		Direct Anthropogenic															
Gas	Units	Net anthropogenic emissions due to Agriculture, Forestry, and Other Land Use (AFOLU)			Non-AFOLU anthropogenic GHG emissions ⁶	Total net anthropogenic emissions (AFOLU + non-AFOLU) by gas			AFOLU as a % of total net anthropogenic emissions, by gas		Natural response of land to human-induced environmental change ⁷			Net land – atmosphere flux from all lands			
Panel 1: Contribution of AFOLU																	
		FOLU		Agriculture	Total												
		A		B	C = A + B		D	E = C + D		F = (C/E) *100		G			A + G		
CO ₂ ²																	
	Gt CO ₂ y ⁻¹	5.2 ± 2.6		No data ¹¹		5.2 ± 2.6		33.9 ± 1.8		39.1 ± 3.2		13%			-11.2 ± 2.6		-6.0 ± 3.7
CH ₄ ^{3,8}	Mt CH ₄ y ⁻¹	19.2 ± 5.8		141.6 ± 42.5		160.8 ± 43		201.3 ± 100.6		362 ± 109							
	Gt CO ₂ e y ⁻¹	0.5 ± 0.2		4.0 ± 1.2		4.5 ± 1.2		5.6 ± 2.8		10.1 ± 3.1		44%					
N ₂ O ^{3,8}	Mt N ₂ O y ⁻¹	0.3 ± 0.1		8.3 ± 2.5		8.7 ± 2.5		2.0 ± 1.0		10.6 ± 2.7							
	Gt CO ₂ e y ⁻¹	0.09 ± 0.03		2.2 ± 0.7		2.3 ± 0.7		0.5 ± 0.3		2.8 ± 0.7		81%					
Total (GHG)	Gt CO ₂ e y ⁻¹	5.8 ± 2.6		6.2 ± 1.4		12.0 ± 2.9		40.0 ± 3.4		52.0 ± 4.5		23%					

Agriculture remains an under represented activity in NDC adaptation and mitigation commitments



Mitigate+: Research
for Low-Emission
Food Systems

50-70% of the countries with the highest potential for reducing GHG emissions in livestock and/or soil carbon included mitigation measures in these subsectors.

- Livestock mitigation priorities include manure management (26 countries), feed management (23 countries) and silvopastoralism (15 countries).
- Soil carbon, mitigation priorities included wetland management (35 countries), agroforestry (34 countries) and grassland management (24 countries).

Many of these activities are also proposed as adaptation measures



Mitigate+: Research
for Low-Emission
Food Systems


Enteric fermentation –Methane



GHG emissions from livestock production vary greatly due to farming practices, animal numbers and type, and food product.



Mitigate+: Research
for Low-Emission
Food Systems



Three promising enteric fermentation mitigating strategies



Mitigate+: Research
for Low-Emission
Food Systems

ENTERIC METHANE MITIGATION STRATEGIES



ANIMAL & FEED MANAGEMENT

- Feed processing
- Genetic selection
- Improving animal health
- Improving pasture management
- Increasing feeding level
- Increasing forage quality
- Optimizing temperature
- TMR feeding

DIET FORMULATION

- By-products
- Decreasing forage-to-concentrate ratios
- Minerals and salts
- Oils and fats
- Oilseeds
- Increasing protein
- Tanniferous forages
- Urea






RUMEN MANIPULATION

- Additives
- Defaunation
- Electron sinks







Effects of mitigation strategies on CH₄ emission

A


Product-Based Reductions


	Mitigation Strategy	Potential Emissions Reduction		Relevant Production System
1	Increasing Feeding Level	CH ₄ I _M CH ₄ I _G	-17% No Data	 
2	Decreasing Grass Maturity	CH ₄ I _M CH ₄ I _G	-13% No Data	 
3	Decreasing Dietary Forage-to-Concentrate Ratio	CH ₄ I _M CH ₄ I _G	-9% -9%	

Absolute Reductions

1	CH ₄ Inhibitors	CH ₄ I _M CH ₄ I _G	-32% No Data	Daily CH ₄ CH ₄ Y	-35% -34%	
2	Tanniferous Forages	CH ₄ I _M CH ₄ I _G	-18% No Data	Daily CH ₄ CH ₄ Y	-12% -10%	 
3	Electron Sinks	CH ₄ I _M CH ₄ I _G	-13% -12%	Daily CH ₄ CH ₄ Y	-17% -15%	
4	Oils & Fats	CH ₄ I _M CH ₄ I _G	-12% -22%	Daily CH ₄ CH ₄ Y	-19% -15%	
5	Oilseeds <small>Lactating animals only</small>	CH ₄ I _M CH ₄ I _G	-12% No Effect	Daily CH ₄ CH ₄ Y	-20% -14%	

Production system

 Feedlot & Mixed Systems

 Grassland Systems

Arndt et al., 2022 Scenario analysis conclusions

Agricultural methane emissions must be decreased by 11 to 30% of the 2010 level by 2030 and by 24 to 47% by 2050 to meet the 1.5 °C target.

Globally, only 100% adoption of the most effective product based and absolute CH₄ reduction strategies can meet the 1.5 °C target by 2030 but not 2050⁴.

Mitigation effects are offset by projected increases in CH₄ due to increasing milk and meat demand.

Notably, by 2030 and 2050, low- and middle-income countries may not meet their contribution to the 1.5 °C target for this same reason, whereas high-income countries could meet their contributions due to only a minor projected increase in enteric CH₄ emissions.



Mitigate+: Research
for Low-Emission
Food Systems

Soils – Nitrous oxide




Regional soil N₂O emissions




Mitigate+: Research
for Low-Emission
Food Systems

N O (Tg)

N_2O emissions are driven by fertilizer application rates and by the levels of N applied in excess of crop demand



Improving N use efficiency and reducing excess N by 75% can reduce N₂O emissions by ~35%





Mitigate+: Research
for Low-Emission
Food Systems

Sinks in agricultural landscapes




There are a wide range of land-based CO₂ removal opportunities



Mitigate+: Research
for Low-Emission
Food Systems


Carbon Dioxide Removal




Soil C sequestration offers significant opportunities across the globe with productivity and soil health benefits



Mitigate+: Research
for Low-Emission
Food Systems




Forests are a key part of the net zero equation



Forests provide climate benefits and buffer against extreme heat events from $\pm 50^\circ\text{N/S}$



Mitigate+: Research
for Low-Emission
Food Systems



Cost-effective (< \$100/tCO₂eq) land-based mitigation is 8–13.8 GtCO₂eq yr⁻¹ between 2020 and 2050



Mitigate+: Research
for Low-Emission
Food Systems


Mitigation category for (a) and (b)

Sectoral:

- Forests and other ecosystems – manage
- Forests and other ecosystems – protect
- Forests and other ecosystems – restore
- Agriculture – reduce emissions
- Agriculture – sequester carbon
- Demand-side


IAM:

- ▨ Forests and other ecosystems – protect (reduce land use change)
- ▨ Forests and other ecosystems – manage and restore (enhance carbon)
- ▨ Agriculture – reduce emissions
- ▨ BECCS




Global Totals (GtCO₂eq year⁻¹)

Sectoral




Technical
32.7 ± 7.2




Cost-effective
13.8 ± 3.1

IAM



IAM
8.0 (0.8 to 16.5)

Total cost-effective potential (GtCO₂eq year⁻¹)



Sectoral

T = Technical
C = Cost-effective
(\$100/tCO₂)

IAM

M = Cost-effective
(\$100/tCO₂)

Roe et al., analyzed feasibility of mitigation measures



Mitigate+: Research
for Low-Emission
Food Systems


The cost-effective potential is approximately 50% from forests and other ecosystems, 35% from agriculture, and 15% from demand-side measures.

Opportunities among countries vary widely depending on types of land-based measures available, their potential co-benefits and risks, and their feasibility.

Governance, economic investment, and socio-cultural conditions influence the likelihood that land-based mitigation potentials are realized.

Assisting countries to overcome barriers may result in significant quantities of near-term, low-cost mitigation while locally achieving important climate adaptation and development benefits.

www.cgiar.org



What is the CGIAR doing to support low emissions food system solutions



The CGIAR is investing ~\$300M per year in improving the sustainability of food systems across developing countries and emerging economies.

2 flagship initiatives focused on climate change

- Mitigate+: Research for low-emission food systems focuses on reducing greenhouse gas emissions from food systems and the predicted consequences of climate change on sustainable development and social equity.
- Climber: This Initiative aims to transform the climate adaptation capacity of food, land and water systems to increase the resilience of smallholder production systems to withstand severe climate change effects like drought, flooding and high temperatures.

Conclusions



Mitigate+: Research
for Low-Emission
Food Systems

Significant GHG emissions reductions in agriculture are achievable through reducing emission intensity of production, but population growth and dietary changes may offset absolute emissions reductions.

Emission reductions consistent with 2030 targets are achievable in many countries; attaining the 2050 targets will require innovation and systems transformation.

Net zero agriculture cannot be achieved without sinks!

Governance, economics, and sociocultural factors are the keys to food system transformations.

Thank you

For discussion:

Emissions reductions are technically feasible

Some countries can meet some targets some of the times

Innovation is needed

Sinks must be part of the solution

Governance, finance, and enabling conditions are the keys to success

Louis Verchot
Alliance Bioversity-CIAT
CGIAR
l.verchot@cgiar.org



Mitigate+: Research
for Low-Emission
Food Systems